


9-CONIC SECTION

Parabola :

The locus of a point which moves such that its distance from a fixed point is equal to its distance from a fixed straight line, i.e. $e = 1$ is called a parabola.



Its equation in standard form is $y^2 = 4ax$

- (i) Focus S (a, 0)
- (ii) Equation of directrix ZM is $x + a = 0$
- (iii) Vertex is O (0, 0)
- (iv) Axis of parabola is X'OX

Some definitions :

Focal distance : The distance of a point on parabola from focus is called focal distance. If P(x_1, y_1) is on the parabola, then focal distance is $x_1 + a$.

Focal chord : The chord of parabola which passes through focus is called focal chord of parabola.

Latus rectum : The chord of parabola which passes through focus and perpendicular to axis of parabola is called latus rectum of parabola. Its length is $4a$ and end points are L(a, 2a) and L'(a, -2a).

Double ordinate : Any chord which is perpendicular to the axis of the parabola is called its double ordinate.

- **Equation of tangent at P(x_1, y_1)** is $yy_1 = 2a(x + x_1)$ and equation of tangent in slope form is $y = mx + \frac{a}{m}$

Here point of contact is $\left(\frac{a}{m^2}, \frac{2a}{m}\right)$

- **Equation of normal at P (x_1, y_1)** is

$$y - y_1 = \frac{-y_1}{2a}(x - x_1)$$

and equation of normal in slope form is $y = mx - 2am - am^3$

Here foot of normal is $(am^2, -2am)$

- The line $y = mx + c$ may be tangent to the parabola if $c = a/m$ and may be normal to the parabola if $c = -2am - am^3$.
- Chord of contact at point (x_1, y_1) is $yy_1 = 2a(x + x_1)$


Ellipse :

If a point moves in a plane in such a way that ratio of its distances from a fixed point (focus) and a fixed straight line (directrix) is always less than 1, i.e. $e < 1$ called an **ellipse**

- Standard equation of an ellipse is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

where $b^2 = a^2(1 - e^2)$

Now, When $a > b$



In this position,

- (i) Major axis $2a$ and minor axis $2b$
- (ii) Foci, S'(-ae, 0) and S(ae, 0) and centre O(0, 0)
- (iii) Vertices A' (-a, 0) and A(a, 0)
- (iv) Equation of directries ZM and Z'M' are

$$x \pm \frac{a}{e} = 0, Z\left(\frac{a}{e}, 0\right) \text{ and } Z'\left(-\frac{a}{e}, 0\right)$$

- (v) Length of latus rectum is $\frac{2b^2}{a} = LL' = L_1L_1'$

- The coordinates of points of intersection of line $y = mx + c$ and the ellipse are given by

$$\left(\frac{-a^2m}{\sqrt{b^2 + a^2m^2}}, \frac{b^2}{\sqrt{b^2 + a^2m^2}}\right)$$

- Equation of tangents of ellipse in term of m is $y = mx \pm \sqrt{b^2 + a^2 m^2}$ and the line $y = mx + c$ is a tangent of the ellipse, if $c = \pm \sqrt{b^2 + a^2 m^2}$
- The length of chord cuts off by the ellipse from the line $y = mx + c$ is

$$\frac{2ab\sqrt{1+m^2} \cdot \sqrt{a^2 m^2 + b^2 - c^2}}{b^2 + a^2 m^2}$$

- The equation of tangent at any point (x_1, y_1) on the ellipse is

$$\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$$

and at the point $(a \cos \phi, b \sin \phi)$ on the ellipse, the tangents is

$$\frac{x \cos \phi}{a} + \frac{y \sin \phi}{b} = 1$$

Parametric equations of the ellipse are

$$x = a \cos \theta \text{ and } y = b \sin \theta.$$

- The equation of normal at any point (x_1, y_1) on the ellipse is

$$\frac{(x - x_1)a^2}{x_1} = \frac{(y - y_1)b^2}{y_1}$$

also at the point $(a \cos \phi, b \sin \phi)$ on the ellipse, the equation of normal is

$$ax \sec \phi - by \operatorname{cosec} \phi = a^2 - b^2$$

- Focal distance of a point $P(x_1, y_1)$ are $a \pm ex_1$
- Chord of contact at point (x_1, y_1) is

$$\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$$

- Chord whose mid-point is (h, k) is

$$\frac{hx}{a^2} + \frac{ky}{b^2} = \frac{h^2}{a^2} + \frac{k^2}{b^2} \text{ i.e. } T = S_1$$

- The locus of point of intersection of two perpendicular tangents drawn on the ellipse is $x^2 + y^2 = a^2 + b^2$. This locus is a circle whose centre is the centre of the ellipse and radius is length of line joining the vertices of major and minor axis. This circle is called "director circle".
- The eccentric angle of point P on the ellipse is made by the major axis with the line PO , where O is centre of the ellipse.
- (a) The sum of the focal distance of any point on an ellipse is equal to the major axis of the ellipse.
- (b) The point (x_1, y_1) lies outside, on or inside the ellipse $f(x, y) = 0$ according as $f(x_1, y_1) > 0$ or < 0 .
- The locus of mid-point of parallel chords of an ellipse is called its **diameter** and its equation is $y = \frac{-b^2 x}{a^2 m}$ which is passes through centre of the ellipse.

- The two diameter of an ellipse each of which bisect the parallel chords of others are called **conjugate diameters**. Therefore, the two diameters $y = m_1 x$ and $y = m_2 x$ will be conjugate diameter if $m_1 m_2 = -\frac{b^2}{a^2}$.

Hyperbola :


When the ratio (defined in parabola and ellipse) is greater than 1, i.e. $e > 1$, then the conic is said to be hyperbola.

Since the equation of the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

differs from that of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ in

having $-b^2$, most of the results proved for the ellipse are true for the hyperbola, if we replace b^2 by $-b^2$ in their proofs. We therefore, give below the list of corresponding results applicable in case of hyperbola.

- Standard equation of hyperbola is $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ where $b^2 = a^2 (e^2 - 1)$



In this case,

- Foci are $S(ae, 0)$ and $S'(-ae, 0)$.
- Equation of directrices ZM and $Z'M'$ are $x \mp \frac{a}{e} = 0, Z\left(\frac{a}{e}, 0\right)$ and $Z'\left(-\frac{a}{e}, 0\right)$
- Transverse axis $AA' = 2a$, conjugate axis $BB' = 2b$.
- Centre $O(0, 0)$.
- Length of latus rectum $LL' = L_1 L_1' = \frac{2b^2}{a}$
- The difference of focal distance from any point $P(x_1, y_1)$ on hyperbola remains constant and is equal to the length of transverse axis. i.e. $S'P - SP = (ex_1 + a) - (ex_1 - a) = 2a$
- The equation of **rectangular hyperbola** $x^2 - y^2 = a^2 = b^2$ i.e. in standard form of hyperbola put $a = b$. Hence $e = \sqrt{2}$ for rectangular hyperbola.